

Wirkungsspektrum Maisherbizide

Produkt oder Mischung

	Bluthirse	Borstenhirse	Hühnerhirse	Einjährige Gräser	Quecken	Raygras	Amarantarten	Ausfallkartoffeln	Ausfallsonnenblumen	Franzosenkraut	Weisser Gänsefuß	Hohlzahn (Gluren)	Knötericharten	Klebern	Melden	Schwarzer Nachtschatten	Ackerwinde	Blacken	Disteln	Schachtelhalm
Banvel 4S							●●	●	●●	●●●●	●	●●	●●	●●	●●●●	●●	●●	●●●●	●●●●	
Callisto	●	●	●●●●	●●	●	●	●●	●●●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●●●●	●●	●	●●●●	●●	●
Callisto + Harmony SX + Successor T	●●	●●	●●●●	●●	●	●●	●●	●●●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●●●●	●●	●	●●●●	●●	●
Callisto + Sitradol SC	●●	●●	●●●●	●●	●	●●	●●	●●●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●●●●	●●	●	●●●●	●●	●
Callisto + Successor T	●●	●●	●●●●	●●		●	●●●●	●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●●●●	●●				●
Callisto + Successor T + Banvel 4S	●●	●●	●●●●	●●	●	●●	●●	●●●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●●●●	●●	●●	●●●●	●●	●
Callisto + Titus + Successor T	●●	●●	●●●●	●●●●	●●	●●	●●	●●●●	●●	●●	●●	●●	●●	●●	●●●●	●●	●	●●	●	●
Principal + Successor T	●●●●	●●●●	●●●●	●●●●	●●●●	●●	●●	●	●●	●●	●●	●●	●●	●●	●●●●	●●	●			
Sitradol SC + Successor T	●●	●●	●●●●	●●		●●	●●		●●	●●	●●	●●	●●	●●	●●●●	●●				
Successor T + Task + Gondor	●●	●●	●●●●	●●●●	●●	●●	●●	●	●●	●●	●●	●●	●●	●●	●●●●	●●	●●	●●●●	●●	
Titus + Exell	●●	●●	●●●●	●●●●	●●	●●	●●		●●	●●	●●	●●	●●	●●	●	●				

● Schwache Wirkung

●● Gute Wirkung bei optimalen Bedingungen

●●● Gute bis sehr gute Wirkung